



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

SEND ALL COMMUNICATIONS ABOUT PROBLEMS TO B. F. FINKEL, Springfield, Mo.

ALGEBRA.

489. Proposed by S. A. COREY, Albia, Iowa.

Prove or disprove the following:

$$\begin{vmatrix} -x & -ay & -bu & abv \\ y & x & -bv & -bu \\ u & av & x & ay \\ -v & -u & y & -x \end{vmatrix}^2 + \begin{vmatrix} a & x & -x & -bu & abv \\ y & y & -bv & -bu \\ u & u & x & ay \\ v & -v & y & -x \end{vmatrix}^2 \\ + \begin{vmatrix} b & x & -ay & -x & abv \\ y & x & y & -bu \\ u & av & u & ay \\ v & -u & -v & -x \end{vmatrix}^2 + \begin{vmatrix} ab & x & -ay & -bu & -x \\ y & x & -bv & y \\ u & av & x & u \\ v & -u & y & -v \end{vmatrix}^2 = \begin{vmatrix} x & -ay & -bu & abv \\ y & x & -bv & -bu \\ u & av & x & ay \\ v & -u & y & -x \end{vmatrix}^2.$$

490. Proposed by HENRY HEATON, Atlantic, Iowa.

Show that $\sin 3^\circ = \frac{1}{16}(\sqrt{30} + \sqrt{10} - \sqrt{6} - \sqrt{2}) + \frac{1}{8}(\sqrt{5} + \sqrt{5} - \sqrt{15} + 3\sqrt{5})$.

491. Proposed by J. W. LASLEY, University of North Carolina.

Solve the equations $xy = x^2 - y^2$ and $x^2 + y^2 = x^3 - y^3$ for x and y .

GEOMETRY.

522. Proposed by GEORGE Y. SOSNOW, Newark, N. J.

Prove that the sum of the squares of the edges of a tetrahedron is equal to four times the sum of the squares of the lines joining the middle points of the opposite edges.

523. Proposed by H. CAMPBELL, St. Johnsbury, Vt.

Given the difference of the segments of the base made by the perpendiculars let fall from the vertical angle, the difference of the base angles, and the sum of the sides of a triangle, to construct the triangle.

CALCULUS.

437. Proposed by LEIGH PAGE, Yale University.

Integrate

$$\int_{-\infty}^{+\infty} \frac{\sin^2 x}{x^2} dx$$

without the use of Gamma Functions.

438. Proposed by PAUL CAPRON, U. S. Naval Academy.

Find the locus of the equation

$$y^6 - 3(a^2 - x^2)y^4 - 2a^2x^2y^3 + 3(a^2 - x^2)y^2 - 6ax^2(a^2 - x^2)y + a^2x^4 - (a^2 - x^2)^3 = 0,$$

first showing that it can be reduced to the form $y = kx^n \pm (a^2 - x^2)^m$, and finding the points of maximum abscissas, of maximum ordinates, and of inflection.

439. Proposed by CLIFFORD N. MILLS, Brookings, South Dakota.

Find the volume of the greatest rectangular parallelopiped that can be inscribed in the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1.$$